

Fresh look on triality

M. Faber

Institut für Kernphysik, Technische Universität Wien, A-1040 Vienna

O. Borisenko, S. Mashkevich and G. Zinovjev

Institute of Theoretical Physics, Ukrainian Academy of Science, Kiev 143

Abstract

Investigating the Z_3 symmetry in Quantum Chromodynamics (QCD) we show that full QCD with a vacuum of vanishing baryonic number does not lead to metastable phases. Rather in QCD with dynamical fermions, the degeneracy of Z_3 phases manifests itself in observables without open triality.

IK-TUW-Preprint 9308401

August 1993

PACS numbers: 12.38.Aw, 12.38Gc

Typeset using REVTeX

The Lagrangian in pure gluonic Quantum Chromodynamics (QCD) has global Z_3 symmetry. It is very well known that in the low temperature phase Z_3 symmetry leads to confinement of color charges and in the high temperature phase Z_3 symmetry is spontaneously broken, color charges are screened by the gluon field. In QCD with dynamical quarks the fermionic contribution to the Lagrangian breaks Z_3 symmetry explicitly.

Recently, in refs. [1,2] it has been concluded on the one hand that this explicit Z_3 symmetry breaking leads to the existence of metastable states at arbitrary high temperatures. On the other hand the authors of refs. [3,4] believe that Z_N phases cannot be prepared as real macroscopic systems. In the following we will try to remove some confusion and show that QCD with dynamical fermions has degenerate Z_3 phases and therefore also ordered-ordered phase transitions like pure gluonic QCD. Furthermore, we emphasize that QCD is able to describe Z_3 symmetric systems only, i.e. systems of zero triality. In order to introduce our concept we study first the pure gluonic system and then turn to QCD with dynamical fermions.

In lattice QCD the gluon field appears in the form of $SU(3)$ matrices $U_{x,\mu}$ which are defined on links (x, μ) of a four-dimensional euclidean lattice. In finite temperature lattice QCD the correlation function $\langle L(\vec{r}_1) \cdots L^*(\vec{r}_N) \rangle$ of several Polyakov loops

$$L(\vec{r}) = \frac{1}{3} \text{Tr} \prod_{t=1}^{N_t} U_{(\vec{r},t),0} \quad (1)$$

can be connected [5] with the free energy F of N infinitely heavy quarks q or antiquarks $\bar{q}$ at the corresponding positions $\vec{r}_1, \dots, \vec{r}_N$ at temperature T relative to the free energy of the vacuum

$$F(q(\vec{r}_1), \dots, \bar{q}(\vec{r}_N)) = -T \ln \langle L(\vec{r}_1) \cdots L^*(\vec{r}_N) \rangle. \quad (2)$$

The thermodynamical average $\langle \cdots \rangle$ is computed using the partition function

$$Z = \int \mathcal{D}[U] e^{-S[U]} \quad (3)$$

as a "sum" over all gauge field configurations U . In pure gluonic QCD for most problems the appropriate choice for S is the Wilson action

$$S_G[U] = \beta \sum_{x, \mu < \nu} \left(1 - \frac{1}{3} \text{Re Tr } U_{x, \mu\nu} \right), \quad \beta = \frac{6}{g^2}, \quad (4)$$

$$U_{x, \mu\nu} = U_{x, \mu} U_{x+\hat{\mu}, \nu} U_{x+\hat{\nu}, \mu}^\dagger U_{x, \nu}^\dagger,$$

where the plaquettes $U_{x, \mu\nu}$ are built of four links in the $\mu\nu$ -plane of a four dimensional euclidean lattice.

The action $S_G[U]$ is Z_3 symmetric. This means that multiplication of all links in direction $\mu = 0$ in the three dimensional x,y,z-torus with fixed t, e.g. $t = 0$ by a Z_3 element leaves the action invariant. A single Polyakov loop $L(\vec{r})$ transforms under Z_3 non-trivially (its triality $\mathcal{T}$ is one), and therefore the distribution of L values can be used as an indicator for spontaneous breaking of Z_3 symmetry.

Let us now discuss the results of lattice calculations and their common interpretation. Monte-Carlo calculations in lattice QCD show usually a characteristic behaviour of the spatial average L of $L(\vec{r})$ during a Monte-Carlo simulation. In the confinement regime (low β) L scatters symmetrically around zero in the complex plane. In the deconfined phase there appear three Z_3 symmetric maxima of the Polyakov loop distribution in 0° - and $\pm 120^\circ$ -directions. As the number of tunneling events between the maxima decreases with increasing β one commonly expects L to be frozen in the thermodynamical limit in one of the Z_3 directions and thus to get spontaneous Z_3 symmetry breaking. The appearance of the three peak structure of the L -distribution in the deconfinement regime is considered as a demonstration that spontaneous symmetry breaking on a finite lattice can never happen exactly. In the thermodynamical limit one may obtain $\arg \langle L \rangle = 1, \pm \frac{2\pi}{3}$. Besides the unpleasant fact that for $\arg(\langle L \rangle) = \pm \frac{2\pi}{3}$ one should reconsider the definition (2) the above interpretation of Monte-Carlo results leads in full QCD to doubtful consequences. Therefore, we prefer another point of view: As $\int \mathcal{D}[U] e^{-S_G[U]}$ is Z_3 invariant and the Polyakov loop (1) is not, one has to get

$$\langle L \rangle = 0 \quad (5)$$

in the confined and also in the deconfined phase. There is no contradiction between this

statement and the spontaneous breaking of Z_3 symmetry in the deconfined phase. It rather shows that the expectation value $\langle L \rangle$ is not the appropriate observable for the demonstration of spontaneous breaking of Z_3 symmetry. One should rather use $\langle L(0)L^*(\vec{r}) \rangle$ correlations or the (Z_3 symmetric) L distributions. The latter is equivalent to the determination of Polyakov loops in Z_3 invariant representations (octet, decuplet, antidecuplet, etc). After this rather trivial observation that for a Z_3 invariant Lagrangian observables of non vanishing triality have expectation value zero we will turn to the more interesting case of full QCD.

In full QCD with dynamical fermions the action S contains a fermionic contribution S_F

$$S = S_G + S_F,$$

$$S_F = \frac{n_F}{4} \sum_{x,x'} \bar{\psi}_x M_{x,x'} \psi_{x'}, M_{x,x'} = D_{x,x'} + m\delta_{x,x'} \quad (6)$$

which breaks Z_3 symmetry explicitly. In the Kogut-Susskind formulation [6] the fermionic matrix M reads

$$M_{x,x'} = \frac{1}{2} \sum_{\mu} \left(\Gamma_{x,\mu} U_{x,\mu} \delta_{x',x+\mu} - \Gamma_{x',\mu} U_{x',\mu}^\dagger \delta_{x,x'-\mu} \right) + m\delta_{x,x'},$$

$$\Gamma_{x,\mu} = (-1)^{x_1+x_2+\dots+x_{\mu-1}}, \quad (7)$$

with one-component Grassmann variables ψ_x and $\bar{\psi}_x$. The integration over the Grassmann variables in the path integral

$$Z = \int \mathcal{D}[U, \bar{\psi}, \psi] e^{-S} \quad (8)$$

can be performed analytically and leads to the fermionic determinant $\det M$

$$Z = \int \mathcal{D}[U] e^{-S_G[U]} \det M. \quad (9)$$

It is easy to see that the fermionic action violates Z_3 symmetry [7]. In the usual MC-iterations one finds that in the low temperature phase the expectation value $\langle L \rangle$ of a Polyakov loop L is a small positive number and in the high temperature phase the three peak structure of the distribution becomes asymmetric. The maximum at 0° is favoured and the two lower maxima at angles of around $\pm 120^\circ$ are symmetric. This fact is usually interpreted as a consequence of the violation of Z_3 symmetry.

Effective models based on this Z_3 asymmetric L -distribution are discussed in refs. [1–4]. As mentioned in the introduction they predict metastable states, but with unphysical properties.

In order to resolve this contradiction we start from the statement that the baryonic number B of the QCD vacuum should be zero. It follows that also the triality $\mathcal{T}$ is zero. Instead of the canonical ensemble with $B = \mathcal{T} = 0$ which should be well suited, a grand canonical description is allowed if it predicts the same behaviour in the thermodynamical limit. Common lattice Monte-Carlo calculations in full QCD use the full fermionic determinant $\det M$ and thus simulate a grand canonical ensemble with chemical potential $\mu = 0$ containing all three triality sectors. There appears now the following loophole in the usual argumentation. The determination of the Polyakov loop expectation value in full QCD gives $\langle L \rangle \neq 0$ and a Z_3 asymmetric L -distribution. In assuming to have measured the L -distribution of the vacuum several authors use effective potentials for the time component of the gluon field arising from one loop effects and predict metastable states [1,2] or find properties of these states [3,4] which are not possible in Minkowski space. However, it has been overlooked that the determinant $\det M$ may be decomposed in three different triality sectors. The decomposition can easily be derived in the following way: Via a hopping parameter expansion the fermionic determinant can be decomposed into contributions according to the number of closed loops of gauge field U which wind around the 4-dimensional lattice in time direction [8–10]. We can then define the triality operator $\hat{\mathcal{T}}$ (with eigenvalues $0, \pm 1$) which determines the triality of such closed loops, and with his help a projection operator $\hat{P}_{\mathcal{T}}$ on triality $\mathcal{T}$

$$\hat{P}_{\mathcal{T}} = \frac{1}{3} \sum_{q=0,\pm 1} e^{q2\pi i(\hat{\mathcal{T}}-\mathcal{T})/3}. \quad (10)$$

Then the triality zero contribution of the fermionic determinant is

$$\det_0 M = \hat{P}_0 \det M = \frac{1}{3} \sum_{q=0,\pm 1} e^{q2\pi i\hat{\mathcal{T}}/3} \det M \quad (11)$$

and more general a projection operator on triality $\mathcal{T}$ §

§The implementation of this projection operator in the lattice formulation is, at least in principle,

$$\hat{P}_{\mathcal{T}} = \frac{1}{3} \sum_{q=0,\pm 1} e^{q2\pi i(\hat{T}-T)/3}. \quad (12)$$

and in general

$$\det M = \sum_{\mathcal{T}=0,\pm 1} \det_{\mathcal{T}} M, \quad (13)$$

with

$$\det_{\mathcal{T}} M = \hat{P}_{\mathcal{T}} \det M. \quad (14)$$

From eq. (13) it is obvious that the observable L having triality $\mathcal{T} = 1$ tests only the $\mathcal{T} = -1$ sector of the fermionic determinant which is not the vacuum sector. Using the correct $\mathcal{T} = 0$ vacuum sector for the fermionic determinant results in a Z_3 symmetric L -distribution, an expectation value $\langle L \rangle = 0$ and no prediction of metastable and unphysical vacuum states. Let us discuss this in more detail. As the measure $\mathcal{D}[U]$ and the gluonic Lagrangian $S_G[U]$ are Z_3 symmetric, the product $\det M[U] O_{\mathcal{T}}[U]$ in

$$\langle O_{\mathcal{T}} \rangle = \frac{1}{Z} \int \mathcal{D}[U] e^{-S_G[U]} \det M[U] O_{\mathcal{T}}[U] \quad (15)$$

has to be also Z_3 symmetric to get $\langle O_{\mathcal{T}} \rangle \neq 0$. In other words Z_3 violating contributions in $\det M[U] O_{\mathcal{T}}[U]$ are automatically eliminated by the path integral. Only the triality $-\mathcal{T}$ component $\det_{-\mathcal{T}} M$ of the fermionic determinant survives. $\det_{-\mathcal{T}} M$ has T quark loops less than antiquark loops winding around the lattice in time direction.

Applying this statement to the example of the single Polyakov loop L representing an infinitely heavy quark Q , the fermionic determinant supplies a triality -1 state of light quarks which will mostly consist of a light antiquark $\bar{q}$ in order to color neutralize the heavy quark Q . A finite value of $\langle L \rangle$ in full QCD therefore does not mean that Z_3 symmetry is

very simple. One has to multiply all links in time direction for an arbitrary time step with the phase $e^{q2\pi i/3}$ and to determine with this new link variables the determinant. The result has to be multiplied with $e^{q2\pi i T/3}$. Finally, the sum over phases $q = 0, \pm 1$ has to be executed.

broken but rather that the fermionic system is flexible enough to neutralize external charges. The Polyakov expectation value $\langle L \rangle$ is a thermodynamical mixture

$$\langle L \rangle = \dots + e^{-F(Q\bar{q})/T} + e^{-F(Qqq)/T} + \dots \quad (16)$$

of a heavy-light meson $Q\bar{q}$, a heavy-light baryon Qqq , This heavy-light states should be bound in the hadronic and "ionized" in the quark-gluon plasma phase. The charge density of such light quarks around static sources has been measured in ref. [11].

For the system of a heavy quark Q at position $\vec{r}_1$ and a heavy antiquark $\bar{Q}$ at position $\vec{r}_2$

$$L(\vec{r}_1)L^*(\vec{r}_2) \quad (17)$$

a totally different component of the fermionic determinant, the triality zero component, contributes. It is evident that this component is Z_3 symmetric, i.e. the three Z_3 transformed sectors which differ in L by factors $e^{\pm\frac{2\pi i}{3}}$ contribute with the same Boltzmann factor. This example shows that for observables of triality zero full QCD is Z_3 symmetric. The fermionic action does not destroy Z_3 symmetry as is the general belief.

We can even extend this statement for triality non zero observables. The above discussion for a single Polyakov line has demonstrated that for Z_3 violating observables full QCD restores the Z_3 symmetry by compensating them to triality zero. Therefore, the effective models of refs. [1–4] are effective models of heavy-light hadrons in the QCD vacuum and not effective models of "the" QCD vacuum.

We would like to summarize that the vacuum of QCD with dynamical fermions has triality zero and therefore degenerate Z_3 phases and ordered-ordered phase transitions like pure gluonic QCD. The Polyakov loop can still be used as an order parameter for the breaking of Z_3 symmetry. There is no need to search new order parameters [12] for full unquenched QCD. In usual lattice Monte-Carlo simulations this triality zero sector can be projected out by using triality zero observables only. Another method would be to use $\det_0 M$ as the fermionic Boltzmann factor. However, an implementation of a projection to fixed triality $\mathcal{T}$ (e.g. $\mathcal{T} = 0$) will be rather complicated due to the fact that usually not $\det M$ is simulated but $(\det M)^2$ whose $\mathcal{T} = 0$ component

$$(\det M)_{\mathcal{T}=0}^2 = (\det_0 M)^2 + 2\det_1 M \det_{-1} M \quad (18)$$

contains contributions from triality ± 1 components. Therefore, one will have to rely on common lattice calculations with the limitations that one should not be misled by the mixing of different triality sectors in the grand canonical ensemble.

M.F. thanks for the warm hospitality at the Institute of Theoretical Physics of the Ukrainian Academy of Science, where the idea of this work came up and to Š. Olejník for critical reading of the manuscript. G.M.Z. wishes to thank J. Engels, B. Petersson and H. Satz for interesting discussions. The work was partially supported by "Fonds zur Förderung der wissenschaftlichen Forschung" under Contract No. P7237-TEC.

After finishing the first version of this article we noticed that J. Polonyi [13] introduced a concept similar to ours. The main conceptual difference to Polonyi's idea is that we demonstrate that common Monte-Carlo calculations use the correct Boltzmann factor for the fermionic vacuum if only triality zero observables are measured.

REFERENCES

- [1] V. Dixit and M. C. Ogilvie, Phys. Lett. B **269**, 353 (1991).
- [2] J. Ignatius, K. Kajantie and K. Rummukainen, Phys. Rev. Lett. **68**, 737 (1992).
- [3] V. M. Belyaev, Ian I. Kogan, G. W. Semenoff and N. Weiss, Phys. Lett. B **277**, 331 (1992).
- [4] W. Chen, M. I. Dobroliubov and G. B. Semenoff, Phys. Rev. D **46**, R1223 (1992).
- [5] L. D. McLerran and B. Svetitsky, Phys. Rev. D **24**, 450 (1981).
- [6] J. B. Kogut, Rev. Mod. Phys. **55**, 755 (1983).
- [7] T. Banks and A. Ukawa, Nucl. Phys. B **225**, 145 (1983).
- [8] N. Weiss, Phys. Rev. D **35**, 2495 (1987).
- [9] I. B. Barbour, C. T. H. Davies and Z. Sabeur, Phys. Lett. B **215**, 567 (1988).
- [10] A. Hasenfratz and D. Touissant, Nucl. Phys. B **371**, 539 (1992).
- [11] W. Bürger, M. Faber, H. Markum and M. Müller, Phys. Rev. D **47**, 3034 (1993).
- [12] C. DeTar and L. D. Mc Lerran, Phys. Lett. B **119**, 171 (1982).
- [13] J. Polonyi, in *Selected Topics in Quark Confinement*, edited by J. Polonyi, F. Csikor, A. Patkós and K. Szlachányi (Eötvös University, Budapest, 1992).